

Flooding of Chaotic Eigenstates into Regular Phase Space Islands

Arnd Bäcker, Roland Ketzmerick, and Alejandro G. Monastera

Institut für Theoretische Physik, Technische Universität Dresden, 01062 Dresden, Germany

(Dated: February 8, 2008)

We introduce a criterion for the existence of regular states in systems with a mixed phase space. If this condition is not fulfilled chaotic eigenstates substantially extend into a regular island. Wave packets started in the chaotic sea progressively flood the island. The extent of flooding by eigenstates and wave packets increases logarithmically with the size of the chaotic sea and the time, respectively. This new effect is observed for the example of island chains with just 10 islands.

PACS numbers: 05.45.Mt, 03.65.Sq

One of the cornerstones in the understanding of the structure of eigenstates in quantum systems is the semiclassical eigenfunction hypothesis [1]: in the semiclassical limit the eigenstates concentrate on those regions in phase space which a typical orbit explores in the long-time limit. For integrable systems these are the invariant tori. For ergodic dynamics the eigenstates become equidistributed on the energy shell [2]. Typical systems have a mixed phase space, where regular islands and chaotic regions coexist. In this case the semiclassical eigenfunction hypothesis implies that the eigenstates can be classified as being either regular or chaotic according to the phase-space region on which they concentrate. Note, that this may fail for an infinite phase space [3].

In this paper we study mixed systems with a compact phase space, but away from the semiclassical limit. Here the properties of eigenstates depend on the size of phase-space structures compared to Planck's constant $\hbar$. In the case of 2D maps this can be very simply stated [4]: a regular state with quantum number $m = 0, 1, \dots$ will concentrate on a torus enclosing an area $(m + 1/2)\hbar$, as can be seen in Fig. 1(c).

We will show that this WKB-type quantization rule is not a sufficient condition. We find a second criterion for the existence of a regular state on the m -th quantized torus,

$$\gamma_m < \frac{1}{\tau_H}. \quad (1)$$

Here $\tau_H = \hbar/\Delta_{\text{ch}}$ is the Heisenberg time of the chaotic sea with mean level spacing Δ_{ch} and γ_m is the decay rate of the regular state m if the chaotic sea were infinite. Quantized tori violating this condition will not support regular states. Instead, chaotic states will *flood* these regions, see Fig. 1(a). In terms of dynamics we find that wave packets started in the chaotic sea progressively flood the island as time evolves. Partial and even complete flooding is possible, depending on system properties. These findings are relevant for islands surrounded by a large chaotic sea.

We numerically demonstrate the flooding and the disappearance of regular states for the important case of island chains. In typical Hamiltonian systems they appear around any regular island. On larger scales they are relevant for Hamiltonian ratchets [5], the kicked rotor with accelerator modes [6], and the experimentally

[7, 8, 9] and theoretically [10] studied kicked atom systems. The flooding of regular islands by chaotic states is a new quantum signature of a classically mixed phase space. This phenomenon shows that not only local phase-space structures, but also global properties of the phase space determine the characteristics of quantum states.

Before we explain the origin of Eq. (1), we numerically study its consequences. We choose a system, where we can change τ_H by increasing the system size without affecting the rates γ_m . A one dimensional kicked system

$$H(p, x, t) = T(p) + V(x) \sum_n \delta(t - n), \quad (2)$$

has a stroboscopic time-evolution given by the mapping, $x_{t+1} = x_t + T'(p_t)$, $p_{t+1} = p_t - V'(x_{t+1})$. The phase space is compact with periodic boundary conditions for $x_t \in [0, M]$ and $p_t \in [-1/2, 1/2]$. Choosing the functions $V'(x)$ and $T'(p)$ appropriately [11] we get a chain of M islands, one per unit cell (see Fig. 1). The islands cover a relative area $A_{\text{reg}} = 0.215$ and have fine structure close to their boundary that is negligible for the quantum proper-




FIG. 1: (a) Husimi representation of a chaotic state flooding the regular islands. Shown are three phase-space cells out of $M = 1597$. Full curves are classical tori close to the border of the regular islands and black dots are iterates of a chaotic orbit. This eigenstate extends well into the islands, while having no weight in their central region. (b) Average of the eigenstate over all M cells. (c) For $M = 1$ eigenstates concentrate either on the chaotic component (left) or over the m -th quantized regular torus. For all plots $\hbar_{\text{eff}} \approx 1/30$ is used.

ties studied here. Points inside an island are mapped one unit cell to the right, i.e. the island chain is transporting.

The eigenstates $|\psi\rangle$ of the quantum system are determined by the eigenvalue equation, $\hat{U}|\psi\rangle = e^{i\varphi}|\psi\rangle$, where $\hat{U}$ is the unitary time-evolution operator over one time period, $\hat{U} = \exp[-2\pi iT(\hat{p})/h_{\text{eff}}] \exp[-2\pi iV(\hat{x})/h_{\text{eff}}]$. The spatial periodicity after M cells requires an effective Planck's constant $h_{\text{eff}} = M/N$, with incommensurate integers M and N . We choose for M/N the rational approximants of $h_{\text{eff}} = 1/(d + \sigma)$ with $\sigma = (\sqrt{5} - 1)/2$ the golden mean and, e.g., $d = 29$ in Fig. 1. This ensures that there are no undesired periodicities and that h_{eff} is approximately constant when varying M . Moreover, the operator $\hat{U}$ reduces to an $N \times N$ periodic band matrix. Using the symmetrized version of the map and making a unitary transformation to a band matrix we are able to calculate the eigenstates $|\psi\rangle$ of $\hat{U}$ up to $N \approx 10^5$.

For $M = 1$, Fig. 1(c) shows a typical chaotic eigenstate and five regular states. The chaotic state extends over the chaotic phase-space component and the regular states concentrate on quantized tori. The eigenstates are represented on the classical phase space by the Husimi distribution, where for visualization we use tilted coherent states adapted to the shape of the island. For larger system sizes we find that chaotic states flood the islands of classically regular motion. Fig. 1(a) shows such a state for $M = 1597$ that clearly ignores the outer tori of the island, which for $M = 1$ act as barriers for chaotic states (Fig. 1(c), left). In the central part of the island, however, this state has essentially no weight. This partial flooding of the island is observed even better in Fig. 1(b), where an average of the Husimi function of this state is taken over all M unit cells. The almost constant value in the chaotic component extends well into the island. Inside the island, clearly away from its outer boundary, the Husimi function sharply drops to zero.

For a quantitative description of this flooding we now analyze the weight W of each eigenstate inside the islands. We determine this weight by integrating the normalized Husimi function (calculated on a 30×30 grid per unit cell) over the islands. In the semiclassical limit, $h_{\text{eff}} \rightarrow 0$, regular states have $W = 1$, while chaotic states have $W = 0$. The distribution $P(W)$ of these weights for all eigenstates is shown in Fig. 2 for various system sizes and $h_{\text{eff}} \approx 1/10$ [12]. For $M = 1$ we observe, as expected, a main peak near $W = 0$ coming from the chaotic eigenstates and two distinct peaks at larger W from the two regular states, see Fig. 2(a). A remarkable shift of the main peak of $P(W)$ to larger values of W can be observed in Fig. 2(b). This shows that by enlarging the system size M all chaotic states continuously increase their weight inside the regular islands. This increase stops when the center of the main peak reaches $W = 0.22$, which corresponds to the area A_{reg} of the island. For these system sizes all states completely flood the island (Fig. 2(c), inset), as observed in Ref. [3].

What happens to the regular states as M is increased? Figure 2(b) shows that the corresponding peaks in the




FIG. 2: Distribution $P(W)$ of the weights W of the eigenstates in the regular islands for $h_{\text{eff}} \approx 1/10$. (a) $M = 1$: The main peak near $W = 0$ is due to chaotic states and two further peaks are due to regular states $m = 0, 1$ (see insets). (b) For increasing M the main peak shifts to larger values of W (white line) and the two peaks from the regular states disappear sequentially. (c) $M = 10946$: A narrow peak remains around $W = A_{\text{reg}}$. Three phase space cells of an eigenstate show a complete flooding of the islands.

distribution $P(W)$ disappear. Notably, the peak for $m = 0$ is much longer visible than the peak for $m = 1$. The nemesis of the regular states can be quantified by determining their fraction f_{reg} as a function of M . To this end we define a state to be regular when its weight W inside the islands exceeds 50%, where the exact criterion does not affect our analysis. Figure 3 shows that the fraction f_{reg} decreases from approximately A_{reg} all the way to zero. The decay is slower for smaller h_{eff} .

Remarkably, in Fig. 3 we see strong signatures of the decrease of f_{reg} already for small system sizes $M \approx 10$. This holds even for small values of h_{eff} , where the complete flooding of the island is numerically not accessible. Similarly, a shift of the main peak in Fig. 2(b) is clearly detectable for such small systems. We thus find that par-




FIG. 3: Fraction f_{reg} of regular states vs. system size M for $h_{\text{eff}} \approx 1/5$ (triangles), $1/10$ (circles), and $1/20$ (squares). An approximately linear decrease with $\ln M$ to $f_{\text{reg}} = 0$ can be seen (lines are a guide to the eye). Already for small system sizes $M \approx 10$ and small h_{eff} a significant reduction of regular states is observed.

tial flooding of regular islands is easily observable.

Why do chaotic states flood the islands of regular motion and why do regular states disappear as the system size is increased? Let us consider a single regular island coupled by tunneling to a chaotic sea. If the chaotic sea is infinite, its states form a continuum. A regular state on the m -th quantized torus has a decay rate γ_m to the continuum [13]. Thus, it is not an eigenstate, but it is dissolved into chaotic states. As a consequence, the chaotic states occupy the m -th quantized torus of the island. If the chaotic sea is finite, but large enough, this decay of the m -th regular state may still take place. The condition for the decay is that during the time $1/\gamma_m$ the discrete chaotic spectrum is not resolved, leading to $1/\gamma_m \ll h/\Delta_{\text{ch}} = \tau_{\text{H}}$ [15]. On the other hand, if the chaotic sea is so small that during the time $1/\gamma_m$ the chaotic spectrum is well resolved, then the regular state m does not decay, yielding Eq. (1). Note, that γ_m increases monotonically with m , as for larger m the m -th torus is closer to the boundary of the island.

The quantized tori of an island can thus be grouped into two classes: (i) The inner tori, $m = 0, \dots, m^* - 1$, where condition (1) is fulfilled and regular states exist. (ii) The outer tori, $m = m^*, \dots, m_{\text{max}} - 1$, where Eq. (1) is violated and which is flooded by chaotic states. Here m_{max} is the number of quantized tori at a given h_{eff} . We find for the fraction f_{reg} of regular states and the weight W_{ch} of chaotic states inside the island:

$$f_{\text{reg}} = A_{\text{reg}} \frac{m^*}{m_{\text{max}}}, \quad W_{\text{ch}} = A_{\text{reg}} \left(1 - \frac{m^*}{m_{\text{max}}}\right). \quad (3)$$

Variation of the system size M in our example allows to change the Heisenberg time $\tau_{\text{H}} \sim M$, while keeping the rates γ_m fixed. Enlarging M leads via Eq. (1) to a decrease of m^* , starting from $m^* = m_{\text{max}} - 1$ all the

way to $m^* = 0$. Together with Eq. (3) this explains Fig. 2(b), where the regular state with $m = 1$ disappears before the $m = 0$ state and the weight W_{ch} grows until $W_{\text{ch}} = A_{\text{reg}}$, where the island is completely flooded, see Fig. 2(c). This also explains the decrease of f_{reg} from A_{reg} to 0, as observed in Fig. 3. This decrease occurs over an exponentially large range in M , due to the roughly exponential dependence of γ_m on m . A quantitative understanding requires a theory for the decay rates γ_m , which is a subject of current research on dynamical tunneling [16, 17]. Note, that in the case of chaos-assisted tunneling the splitting of symmetry related regular states fluctuates strongly, depending on individual chaotic states. In contrast, the decay rate γ_m describes an average tunneling to a continuum of chaotic states.

Variation of h_{eff} affects both γ_m and τ_{H} in Eq. (1). While $\tau_{\text{H}} \sim M/h_{\text{eff}}$, one expects in analogy to WKB theory that $\gamma_m \sim \exp[-g(m/m_{\text{max}})/h_{\text{eff}}]$, where the system specific function g decreases monotonically to $g(1) = 0$. From the definition of m^* follows $m^*/m_{\text{max}} = g^{-1}[h_{\text{eff}} \ln(M/h_{\text{eff}})]$, where g^{-1} decreases monotonically. Decreasing h_{eff} reduces the argument of g^{-1} such that m^*/m_{max} increases. Eq. (3) implies that f_{reg} grows and W_{ch} decreases. Note that in the semiclassical limit, $h_{\text{eff}} \rightarrow 0$, we obtain $m^*/m_{\text{max}} \rightarrow 1$. This is in agreement with the semiclassical eigenfunction hypothesis, namely $f_{\text{reg}} = A_{\text{reg}}$ and there is no flooding. In contrast, if the system size is infinite, we have an infinite τ_{H} and our argument leads to $m^* = 0$, i.e. complete flooding, for any h_{eff} . This coincides with the considerations of [3] implying a failure of the semiclassical eigenfunction hypothesis.

Our explanation is complete for systems without localization. For example, this is the case if the average classical drift of a unit cell is non-zero, like in atom optic experiments in the presence of gravity [9, 10]. Localization, however, sets a lower bound to the effective mean level spacing, $\Delta_{\text{ch}} \sim 1/\lambda$, where λ is the localization length. For $M > \lambda$, this leads to $\tau_{\text{H}} \sim \lambda$ and m^* stays at its value for $M = \lambda$. According to Eq. (3) the same holds for f_{reg} and W_{ch} . This applies, e.g., to dynamical localization in the kicked rotor. For transporting islands, like in the model studied here, $\lambda \sim 1/\gamma_0$ is unusually large [3, 14, 18], such that already for $M = \lambda$ one has $m^* = 0$, $f_{\text{reg}} = 0$, and $W_{\text{ch}} = A_{\text{reg}}$. In this case, localization has no consequences [19].

We generalize our analysis to the dynamics of wave packets, which is experimentally of great relevance [9]. A wave packet started on the m -th torus will be restricted to that region if condition (1) is fulfilled, i.e. $m < m^*$. If $m > m^*$, however, the wave packet will decay into the chaotic sea. Particularly interesting is the case of a wave packet started inside the chaotic sea. The island is progressively flooded, i.e. the m -th torus at time $t_m \sim 1/\gamma_m$ for $m > m^*$. For $t > \tau_{\text{H}}$ the weight $W(t)$ will saturate at W_{ch} , Eq. (3). This is confirmed in Fig. 4, for increasing values of M .

Our results have consequences for spectral statistics in mixed systems which go well beyond the previously stud-




FIG. 4: Weight W in the island vs. time t for a wave packet initially started as a coherent state in the chaotic sea at $(x, p) = (0.4, -0.2)$ for system sizes $M = 1, 13, 144, 1597$ and $h_{\text{eff}} \approx 1/30$. The thick (green) line guides the eye to the linear increase with $\ln t$ until $W = A_{\text{reg}}$. Insets show the time-evolved wave packet averaged over all cells for the case $M = 1597$, demonstrating the progressive flooding of the island.

ied effects of dynamical tunneling (see e.g. [16, 20]). The effective size of the regular region, f_{reg} in Eq. (3), entering the Berry-Robnik formula [21] is drastically reduced.

Our analysis applies as well to hierarchical states [22], which are confined by partial transport barriers with turnstile areas smaller than h . We predict the additional condition $\gamma < 1/\tau_H$ for their existence, where γ describes the decay through these partial barriers. For regular states on island chains within that hierarchical region, condition (1) applies, with τ_H given by the mean level spacing of the surrounding hierarchical states.

Finally, we emphasize that the time periodicity of the system (2) and the restriction of our discussion to maps is not crucial and that we expect flooding of islands for any Hamiltonian with a mixed phase space. We stress that this new quantum signature of chaos for eigenstates and wave packet dynamics already appears for small system sizes, e.g. island chains of length 10. This makes numerical explorations very feasible and should lead to experimental observations, for example using optical lattices [7, 8, 9].

We thank Holger Schanz for discussions and the DFG for support under contract KE 537/3-2.

-
- [1] I. C. Percival, J. Phys. B **6**, L229 (1973); M. V. Berry, J. Phys. A **10**, 2083 (1977); A. Voros, in *Stochastic Behavior in Classical and Quantum Hamiltonian Systems* (Springer, Berlin, 1979), pp. 326.
 - [2] See e.g. A. Bäcker, R. Schubert and P. Stifter, Phys. Rev. E **57**, 5425 (1998), and references therein.
 - [3] L. Hufnagel, R. Ketzmerick, M.-F. Otto, and H. Schanz, Phys. Rev. Lett. **89**, 154101 (2002).
 - [4] M. V. Berry, N. L. Balazs, M. Tabor, and A. Voros, Ann. Phys. **122**, 26 (1979).
 - [5] H. Schanz, M.-F. Otto, R. Ketzmerick, and T. Dittrich, Phys. Rev. Lett. **87**, 070601 (2001).
 - [6] F. M. Izrailev, Phys. Rep. **196**, 299 (1990).
 - [7] D. A. Steck, W. H. Oskay, and M. G. Raizen, Science **293**, 274 (2001).
 - [8] W. K. Hensinger *et al.*, Nature **412**, 52 (2001).
 - [9] M. K. Oberthaler *et al.*, Phys. Rev. Lett. **83**, 4447 (1999); S. Schlunk, M. B. d'Arcy, S. A. Gardiner, and G. S. Summy, Phys. Rev. Lett. **90**, 124102 (2003).
 - [10] S. Fishman, I. Guarneri, and L. Rebuzzini, Phys. Rev. Lett. **89**, 084101 (2002), J. Stat. Phys. **110**, 911 (2003); M. B. d'Arcy *et al.*, Phys. Rev. E **69**, 027201 (2004).
 - [11] We start with linear functions $t'(p) = 1/2 \pm (1 - 2p)$ for $0 < \pm p < 1/2$ (periodically extended) and $v'(x) = -k - r(x - k)$ for $k - 1/2 < x < k + 1/2$, where $k \in \mathbb{Z}$. Smoothing with a Gaussian, $G(z) = \exp(-z^2/2\varepsilon^2)/\sqrt{2\pi\varepsilon^2}$, gives analytic functions $T'(p) = \int dz t'(p + z) G(z)$ and $V'(x) = \int dz v'(x + z) G(z)$. We use $r = 0.65$, and $\varepsilon = 0.015$, as for Fig. 1a in [3].
 - [12] We improve the statistics for small M by varying the phase of the boundary condition in the x -direction, which does not affect the underlying classical dynamics. For large M we use 2000 eigenstates.
 - [13] This is in analogy to a discrete state coupled to a continuum discussed in [U. Fano, Nuovo Cimento, N. s. **12**, 154 (1935)]. In practice one can determine the rates γ_m by starting suitable wave packets as in [14].
 - [14] J. D. Hanson, E. Ott, and T. M. Antonsen, Phys. Rev. A **29**, 819 (1984).
 - [15] A barrier tunneling analogy is given in E. J. Heller and M. J. Davies, J. Phys. Chem **85**, 307 (1981). For a discussion of time scales in quantum chaos see e.g. [F. Borgonovi and G. Casati, in *Frontiers in Quantum Physics* (Springer, Singapore, 1998), pp. 127, cond-mat/9711281]. By Fermi's golden rule one can show that the reasoning using time scales is equivalent to the comparison of the coupling matrix element v_m to the mean spacing Δ_{ch} .
 - [16] V. A. Podolskiy and E. E. Narimanov, Phys. Rev. Lett. **91**, 263601 (2003).
 - [17] M. J. Davis and E. J. Heller, J. Chem. Phys. **75**, 246 (1981); O. Bohigas, S. Tomsovic, and D. Ullmo, Phys. Rep. **223**, 43 (1993); O. Brodier, P. Schlagheck, and D. Ullmo, Ann. Phys. **300**, 88 (2002); C. Eltschka and P. Schlagheck, Phys. Rev. Lett. **94**, 014101 (2005).
 - [18] A. Iomin, S. Fishman, and G. M. Zaslavsky, Phys. Rev. E **65**, 036215 (2002) and references therein.
 - [19] Depending on the factor between λ and $1/\gamma_0$, the $m = 0$ state might couple to very few chaotic states only. The emerging eigenstates may look more like hybrid states that are regular in some unit cells and chaotic in others, explaining such an observation in Fig. 4 of Ref. [3].
 - [20] T. Prosen and M. Robnik, J. Phys. A **27**, 8059 (1994).
 - [21] M. V. Berry and M. Robnik, J. Phys. A **17**, 2413 (1984).
 - [22] R. Ketzmerick, L. Hufnagel, F. Steinbach, and M. Weiss, Phys. Rev. Lett. **85**, 1214 (2000).